

Latest Results on g_1 and g_2 at high x

Jian-ping Chen

Jefferson Lab, Newport News, Virginia 23606, USA

Abstract. Recent progress from Jefferson Lab has significantly improved our understanding of the nucleon spin structure in the high- x region. Results of a precision measurement of the neutron spin asymmetry, A_1^n , in the high- x (valence quark) region are discussed. The up and down quark spin distributions in the nucleon were extracted. A_2^n was also measured. The results were used, in combination with existing data, to extract the second moment, d_2^n . Preliminary results on A_1^p and A_1^d in the high- x region have also become available. Finally, the results of a precision measurement of the g_2 structure function to study higher twist effects will be presented.

Keywords: Spin Structure, high x , higher twist, JLab

PACS: 13.60.Hb

Introduction and Motivation

Recently, the high polarized luminosity available at Jefferson Lab (JLab) has allowed the study of the nucleon spin structure with an unprecedented precision, enabling us to access the valence quark (high- x) region and also to expand the study to the second spin structure function, g_2 .

The high- x region is of special interest, because this is where the valence quark contributions are expected to dominate. With sea quark and explicit gluon contributions expected not to be important, it is a clean region to test our understanding of nucleon structure. Relativistic constituent quark models [1] should be applicable in this region and perturbative QCD [2] can be used to make predictions in the large x limit.

To first approximation, the constituent quarks in the nucleon are described by the SU(6) wavefunctions. SU(6) symmetry leads to the following predictions:

$$A_1^p = 5/9; A_1^n = 0; \Delta u/u = 2/3; \Delta d/d = -1/3. \quad (1)$$

Relativistic Constituent Quark Models (RCQM) with broken SU(6) symmetry, e.g., the hyperfine interaction model [1], lead to a dominance of a ‘diquark’ configuration with the diquark spin $S = 0$ at high x . This implies that as $x \rightarrow 1$:

$$A_1^p \rightarrow 1; A_1^n \rightarrow 1; \Delta u/u \rightarrow 1; \text{ and } \Delta d/d \rightarrow -1/3. \quad (2)$$

In the RCQM, relativistic effects give rise to the quark orbital angular momentum and reduce the valence quark contributions to the nucleon spin from 1 to $0.6 - 0.75$.

Another approach is leading-order pQCD [2], which assumes the quark orbital angular momentum to be negligible and leads to hadron helicity conservation. It yields:

$$A_1^p \rightarrow 1; A_1^n \rightarrow 1; \Delta u/u \rightarrow 1; \text{ and } \Delta d/d \rightarrow 1. \quad (3)$$

Not only are the limiting values as $x \rightarrow 1$ important, but also the behavior in the high- x region. How A_1^n and A_1^p approach their limiting values when x approaches 1, is sensitive to the dynamics in the valence quark region.

g_2 , unlike g_1 and F_1 , cannot be interpreted in the simple quark-parton model. To understand g_2 properly, it is best to start with the operator product expansion method (OPE) [3]. In the OPE, neglecting quark masses, g_2 can be cleanly separated into a twist-2 and a higher twist term:

$$g_2(x, Q^2) = g_2^{WW}(x, Q^2) + g_2^{H.T.}(x, Q^2) . \quad (4)$$

The leading-twist term can be determined from g_1 as [4]

$$g_2^{WW}(x, Q^2) = -g_1(x, Q^2) + \int_x^1 \frac{g_1(y, Q^2)}{y} dy , \quad (5)$$

and the higher-twist term arises from the quark-gluon correlations. Therefore g_2 provides a clean way to study higher-twist effects. In addition, at high Q^2 , the x^2 -weighted moment, d_2 , is a twist-3 matrix element and is related to the color polarizabilities [5]:

$$d_2 = \int_0^1 x^2 [g_2(x) - g_2^{WW}(x)] dx. \quad (6)$$

Predictions for d_2 exist from various models and lattice QCD.

Recent results from Jefferson Lab

In 2001, JLab experiment E99-117 [6] was carried out in Hall A to measure A_1^n with high precision in the x region from 0.33 to 0.61 (Q^2 from 2.7 to 4.8 GeV²). Asymmetries from inclusive scattering of a highly polarized 5.7 GeV electron beam on a high pressure (> 10 atm) (both longitudinally and transversely) polarized ³He target were measured. Parallel and perpendicular asymmetries were extracted for ³He. After taking into account the beam and target polarization and the dilution factor, they were combined to form A_1^{3He} . Using the most recent model [7], nuclear corrections were applied to extract A_1^n . The results on A_1^n are shown in the left panel of Fig. 1.

The experiment greatly improved the precision of data in the high- x region, providing the first evidence that A_1^n becomes positive at large x , showing clear SU(6) symmetry breaking. The results are in good agreement with the LSS 2001 pQCD fit to previous world data [8] (solid curve) and the statistical model [9] (long-dashed curve). The trend of the data is consistent with the RCQM predictions (the shaded band). The data disagree with the predictions from the leading-order pQCD models (short-dashed and dash-dotted curves).

In the leading-order approximation, the polarized quark distribution functions $\Delta u/u$ and $\Delta d/d$ were extracted from our neutron data combined with the world proton data. The results are shown in the right panel of Fig. 1, along with predictions from the RCQM (dot-dashed curves), leading-order pQCD (short-dashed curves), the LSS 2001




FIGURE 1. A_1^n , $\Delta u/u$ and $\Delta d/d$ results compared with the world data and theoretical predictions.

fits (solid curves) and the statistical model (long-dashed curves). The results agree well with RCQM predictions as well as the LSS 2001 fits and statistical model but are in significant disagreement with the predictions from the leading-order pQCD models assuming hadron helicity conservation. This suggests that effects beyond leading-order pQCD, such as the quark orbital angular momentum, may play an important role in this kinematic region.

A_2^n was also obtained from the same experiment. The precision of the A_2^n data is comparable to that of the best existing world data [10] at high x . Combining these results with the world data, the second moment d_2^n was extracted at an average Q^2 of 5 GeV²:

$$d_2^n = 0.0062 \pm 0.0028. \quad (7)$$

Compared to the previously published result [10], the uncertainty on d_2^n has been improved by about a factor of 2. The d_2 moment at high Q^2 has been calculated by Lattice QCD and a number of theoretical models. While a negative or near-zero value was predicted by Lattice QCD and most models, the new result for d_2^n is positive.

Preliminary results of A_1^p and A_1^d from the Hall B eg1 experiment [11] have recently become available. The data cover the Q^2 range of 1.4 to 4.5 GeV² for x from 0.2 to 0.6 with an invariant mass larger than 2 GeV. The precision of the data improved significantly over that of the existing world data.

A precision measurement of g_2^n from JLab E97-103 [12] covered five different Q^2 values from 0.58 to 1.36 GeV² at $x \approx 0.2$. Results for g_2^n as well as g_1^n are given in Fig. 2. The light-shaded area in the two plots gives the leading-twist contribution to these two quantities, respectively, obtained by fitting world data and evolving to the Q^2 values

of this experiment. The systematic errors are shown as the dark-shaded area near the horizontal axes.




FIGURE 2. Fig. 1: results for g_1^n (left) and g_2^n (right) from E97103.

The precision reached is more than an order of magnitude improvement over that of the best world data. The difference of g_2 from the leading twist part (g_2^{WW}) [4] is due to higher twist effects and is sensitive to quark-gluon correlations. The g_2^{WW} values were obtained from a fit [13] to the world high Q^2 data, then evolved to the Q^2 values of this experiment. The measured g_2^n values are consistently higher than g_2^{WW} . For the first time, there is a clear indication that higher twist effects become important at the level of precision of these data. The new g_1^n data agree with the leading-twist calculations within the uncertainties.

In summary, the high polarized luminosity available at JLab, has provided high-precision data to study the nucleon spin structure in the high- x region and higher twist effects, which shed light on the valence quark structure and help to understand quark-gluon correlations.

The work presented was supported in part by the U. S. Department of Energy (DOE) contract DE-AC05-84ER40150 Modification NO. M175, under which the Southeastern Universities Research Association operates the Thomas Jefferson National Accelerator Facility.

REFERENCES

1. N. Isgur, Phys. Rev. D **59**, 034013 (1999).
2. S. Brodsky, M Burkhardt and I. Schmidt, Nucl. Phys. B **441**, 197 (1995).
3. K. Wilson, Phys. Rev. **179**, 1499 (1969).
4. S. Wandzura and F. Wilczek, Phys. Lett. B **72** (1977).
5. X. Ji and P. Unrau, Phys. Lett. B **333**, 228 (1994).
6. X. Zheng, *et al.*, Phys. Rev. Lett. **92**, 012004 (2004); Phys. Rev. C **70**, 065207 (2004).
7. F. Bissey, *et al.*, Phys. Rev. C **65**, 064317 (2002).
8. E. Leader, A. V. Sidorov and D. B. Stamenov, Eur. Phys. J. C **23**, 479 (2002).
9. C. Bourrely, J. Soffer and F. Buccella, Eur. Phys. J. C **23**, 487 (2002).
10. K. Abe, *et al.*, E155 collaboration, Phys. Lett. B **493**, 19 (2000).
11. S. Kuhn, private communication.
12. K. Kramer, *et al.*, nucl-ex/0506005, submitted to Phys. Rev. Lett.
13. J. Blümlein and H. Bottcher, Nucl. Phys. B **636**, 225 (2002).